

Forage yield of wheat and triticale varieties, Stephenville, TX, 1997-98.

Variety	Yield/Acre (lbs DM)				
	Harvest 1 12/11/97	Harvest 2 2/11/98	Harvest 3 3/25/98	Harvest 4 5/12/98	Total
Lockett (wheat)	201 a <sup>1</sup>	607 a	1633 a	531 a	2958 c
TAM-202 (wheat)	171 a	825 a	1775 a	566 a	3337 bc
Jagger (wheat)	313 a	759 a	1908 a	454 a	3784 bc
Custer (wheat)	286 a	566 a	1757 a	515 a	3292 bc
Presto (triticale)	260 a	697 a	2653 a	751 a	4675 a
Hartman (triticale)	214 a	538 a	2057 a	721 a	2986 c
Trit II (triticale)	162 a	260 a	1974 a	890 a	3911 ab
Triplecale (triticale)	188 a	587 a	1834 a	966 a	3576 bc
<b>CV</b>	<b>51</b>	<b>54</b>	<b>23</b>	<b>46</b>	<b>17</b>

<sup>1</sup>Means in the same column with the same letter are not significantly different according to DMRT (P = 0.10).

Table 2. Silage yield and quality of wheat and triticale varieties, Stephenville, TX, 1997-98.

Variety	Harvest Date	Yield/A <sup>1</sup> (tons @ 35% DM)	Harvest Moisture (%)	Crude Protein (%)	ADF (%)	NDF (%)
Lockett (wheat)	4-28-98	11.3 a <sup>2</sup>	54.3 a	6.5 a	29.8 e	54.4 e
TAM-202 (wheat)	4-24-98	9.9 ab	52.5 ab	6.6 a	30.3 de	55.9 de
Jagger (wheat)	4-24-98	8.3 b	52.4 ab	7.0 a	30.7 de	55.9 de
Custer (wheat)	4-24-98	10.6 ab	52.6 ab	6.9 a	32.5 cd	57.6 cd
Presto (triticale)	5-01-98	11.7 a	51.9 ab	7.4 a	32.0 de	57.9 cd
Hartman (triticale)	5-11-98	9.9 ab	47.6 cd	6.6 a	37.1 a	66.4 a
Trit II (triticale)	5-11-98	8.4 b	50.1 bc	7.2 a	35.2 ab	62.9 b
Triplecale (triticale)	5-12-98	11.7 a	46.0 d	6.9 a	34.2 bc	60.3 bc
<b>CV</b>		<b>17.4</b>	<b>4.2</b>	<b>13.9</b>	<b>5.1</b>	<b>3.7</b>

<sup>1</sup>All varieties were harvested in the soft dough stage.

<sup>2</sup>Means in the same column with the same letter are not significantly different according to DMRT (P = 0.10).